


Circular Motion 1

Q1.



A particle P of mass 0.3 kg is attached to a fixed point A by a light inextensible string of length 0.8 m . The fixed point O is 0.15 m vertically below A . The particle P moves with constant speed $v \text{ m s}^{-1}$ in a horizontal circle with centre O (see diagram).


- (i) Show that the tension in the string is 16 N . [2]
- (ii) Find the value of v . [3]
-

Q2.

A particle P of mass 0.4 kg is attached to a fixed point A by a light inextensible string of length 0.5 m . The point A is 0.3 m above a smooth horizontal surface. The particle P moves in a horizontal circle on the surface with constant angular speed 5 rad s^{-1} .

- (i) Calculate the tension in the string. [3]
- (ii) Find the magnitude of the force exerted by the surface on P . [2]
-

Q3.



A light inextensible string of length a is threaded through a fixed smooth ring R . One end of the string is attached to a particle A of mass $3m$. The other end of the string is attached to a particle B of mass m . The particle A hangs in equilibrium at a distance x vertically below the ring. The angle between AR and BR is θ (see diagram). The particle B moves in a horizontal circle with constant angular speed $2\sqrt{\frac{g}{a}}$.

Show that $\cos \theta = \frac{1}{3}$ and find x in terms of a . [5]


Circular Motion 1

Q4.

A particle P of mass m is attached to one end of a light inextensible string of length a . The other end of the string is attached to a fixed point O on a smooth horizontal plane. The particle P moves in horizontal circles about O . The tension in the string is $4mg$.

Find, in terms of a and g , the time that P takes to make one complete revolution. [2]

Q5.



A particle P is attached to one end of a light inextensible string of length a . The other end of the string is attached to a fixed point O . The particle P is held with the string taut and making an angle θ with the downward vertical. The particle P is then projected with speed $\frac{4}{5}\sqrt{5ag}$ perpendicular to the string and just completes a vertical circle (see diagram).

Find the value of $\cos\theta$. [5]

Q6.

A particle P of mass m is moving in a horizontal circle with angular speed ω on the smooth inner surface of a hemispherical shell of radius r . The angle between the vertical and the normal reaction of the surface on P is θ .

(a) Show that $\cos\theta = \frac{g}{\omega^2 r}$. [3]

The plane of the circular motion is at a height x above the lowest point of the shell. When the angular speed is doubled, the plane of the motion is at a height $4x$ above the lowest point of the shell.

(b) Find x in terms of r . [4]
