


Q1.



The diagram shows a semicircle  $ACB$  with centre  $O$  and radius  $r$ . The angle  $BOC$  is  $x$  radians. The area of the shaded segment is a quarter of the area of the semicircle.

(i) Show that  $x$  satisfies the equation

$$x = \frac{3}{4}\pi - \sin x. \quad [3]$$


(ii) This equation has one root. Verify by calculation that the root lies between 1.3 and 1.5. [2]

(iii) Use the iterative formula

$$x_{n+1} = \frac{3}{4}\pi - \sin x_n$$

to determine the root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

Q2.



The diagram shows the curve  $y = \frac{\sin x}{x}$  for  $0 < x \leq 2\pi$ , and its minimum point  $M$ .

(i) Show that the  $x$ -coordinate of  $M$  satisfies the equation

$$x = \tan x. \quad [4]$$

(ii) The iterative formula

$$x_{n+1} = \tan^{-1}(x_n) + \pi$$

can be used to determine the  $x$ -coordinate of  $M$ . Use this formula to determine the  $x$ -coordinate of  $M$  correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

9709/32/M/J/10 Q4

Q3.

The curve  $y = \frac{\ln x}{x+1}$  has one stationary point.

(i) Show that the  $x$ -coordinate of this point satisfies the equation

$$x = \frac{x+1}{\ln x},$$

and that this  $x$ -coordinate lies between 3 and 4. [5]

(ii) Use the iterative formula

$$x_{n+1} = \frac{x_n + 1}{\ln x_n}$$

to determine the  $x$ -coordinate correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

9709/33/M/J/10 Q6

Q4.

- (i) By sketching suitable graphs, show that the equation

$$4x^2 - 1 = \cot x$$

has only one root in the interval  $0 < x < \frac{1}{2}\pi$ . [2]

- (ii) Verify by calculation that this root lies between 0.6 and 1. [2]

- (iii) Use the iterative formula

$$x_{n+1} = \frac{1}{2}\sqrt{1 + \cot x_n}$$

to determine the root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

9709/31/O/N/10 Q4


Q5.

- (i) Given that  $\int_1^a \frac{\ln x}{x^2} dx = \frac{2}{5}$ , show that  $a = \frac{5}{3}(1 + \ln a)$ . [5]

- (ii) Use an iteration formula based on the equation  $a = \frac{5}{3}(1 + \ln a)$  to find the value of  $a$  correct to 2 decimal places. Use an initial value of 4 and give the result of each iteration to 4 decimal places. [3]

9709/33/O/N/1 Q7

Q6.




The diagram shows a circle with centre  $O$  and radius 10 cm. The chord  $AB$  divides the circle into two regions whose areas are in the ratio 1 : 4 and it is required to find the length of  $AB$ . The angle  $AOB$  is  $\theta$  radians.

- (i) Show that  $\theta = \frac{2}{5}\pi + \sin \theta$ . [3]

- (ii) Showing all your working, use an iterative formula, based on the equation in part (i), with an initial value of 2.1, to find  $\theta$  correct to 2 decimal places. Hence find the length of  $AB$  in centimetres correct to 1 decimal place. [5]

9709/31/M/J/11 Q6

Q7.



The diagram shows a semicircle  $ACB$  with centre  $O$  and radius  $r$ . The tangent at  $C$  meets  $AB$  produced at  $T$ . The angle  $BOC$  is  $x$  radians. The area of the shaded region is equal to the area of the semicircle.

(i) Show that  $x$  satisfies the equation

$$\tan x = x + \pi. \quad [3]$$

(ii) Use the iterative formula  $x_{n+1} = \tan^{-1}(x_n + \pi)$  to determine  $x$  correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

9709/32/M/J/11 Q4

Q8.

(i) By sketching a suitable pair of graphs, show that the equation

$$\cot x = 1 + x^2,$$

where  $x$  is in radians, has only one root in the interval  $0 < x < \frac{1}{2}\pi$ . [2]

(ii) Verify by calculation that this root lies between 0.5 and 0.8. [2]

(iii) Use the iterative formula

$$x_{n+1} = \tan^{-1}\left(\frac{1}{1+x_n^2}\right)$$

to determine this root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

9709/33/M/J/11 Q6

## Numerical solutions of equations 1



Q9.

- (i) By sketching a suitable pair of graphs, show that the equation

$$\sec x = 3 - \frac{1}{2}x^2,$$

where  $x$  is in radians, has a root in the interval  $0 < x < \frac{1}{2}\pi$ . [2]

- (ii) Verify by calculation that this root lies between 1 and 1.4. [2]

- (iii) Show that this root also satisfies the equation

$$x = \cos^{-1}\left(\frac{2}{6-x^2}\right). \quad [1]$$

- (iv) Use an iterative formula based on the equation in part (iii) to determine the root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

9709/31/O/N/11 Q5

Q10.

It is given that  $\int_1^a x \ln x \, dx = 22$ , where  $a$  is a constant greater than 1.

- (i) Show that  $a = \sqrt{\left(\frac{87}{2 \ln a - 1}\right)}$ . [5]

- (ii) Use an iterative formula based on the equation in part (i) to find the value of  $a$  correct to 2 decimal places. Use an initial value of 6 and give the result of each iteration to 4 decimal places. [3]

9709/33/O/N/11 Q5